

Effects of Herbal Bath “HAC” on Functional Recovery and c-Fos Expression in the Ventrolateral Periaqueductal Gray Region after Sciatic Crushed Nerve Injury in Rats

Moon-Sang Ryu¹, Hyung-Ho Lim¹, Yun-Kyung Song¹,
Hye-Jung Lee², Jin-Hee Seo³, Myoung-Hwa Lee³,
Mal-Soon Shin³ and Chang-Ju Kim^{3*}

¹Department of Rehabilitation Medicine, College of Oriental Medicine, Kyungwon University, Sungnam 461-701, ²Acupuncture and Meridian Science Research Center, ³Department of Physiology, College of Medicine, Kyung Hee University, Seoul 130-701, Korea

ABSTRACT

Peripheral nerve injuries are a commonly encountered clinical problem and often result in a chronic pain and severe functional deficits. c-Fos expression is sometimes used as a marker of increased neuronal activity. We have developed herbal bath “HAC” for pain control using the following herbs: *Harpagophytum procumbens*, *Atractylodes japonica*, and *Corydalis tuber*. In the present study, we investigated the effects of herbal bath “HAC” on the recovery rate of the locomotor function and the expression of c-Fos in the ventrolateral periaqueductal gray (vlPAG) region of brain following sciatic crushed nerve injury in rats. Walking track analysis for the evaluation of functional recovery and immunohistochemistry for the c-Fos expression were used for this study. In the present results, characteristic gait change with dropping of the sciatic function index (SFI) was observed and c-Fos expression in the vlPAG was suppressed following sciatic crushed nerve injury in rats. Immersion into herbal bath “HAC” enhanced SFI value and restored c-Fos expression in the vlPAG to the control value. These results suggest that herbal bath “HAC” might activate neurons in the vlPAG, and it facilitates functional recovery from peripheral nerve injury. Here we showed that herbal bath “HAC” could be used as a new therapeutic intervention for pain control and functional recovery from peripheral nerve injury.

Key words: herbal bath, sciatic crushed nerve injury, sciatic function index, c-Fos, ventrolateral periaqueductal gray

INTRODUCTION

Crush injury on the sciatic nerve serves as the animal model of unilateral peripheral neuropathy. The affected limb displays characteristics of painful

*To whom correspondence should be addressed.
TEL: 82-2-961-0407, FAX: 82-2-964-2195
e-mail: changju@khu.ac.kr
Received December 10, 2009
Accepted for publication December 21, 2009

neuropathy such as hyperalgesia, pain-related gait, and swelling (Bennett and Xie, 1988). These features are considered as the abnormal responses to peripheral stimuli, reflecting the changes in central nervous system (CNS) nociceptive neural transmission.

The mammalian nervous system contains networks that modulate nociceptive transmission. Of these, the descending pain control system consists of three major components: the periaqueductal gray (PAG) of the midbrain, the rostroventral medulla (RVM) including the nucleus raphe magnus (NRM), and the spinal dorsal horn. Neurons in the PAG and NRM project directly to the spinal cord dorsal horn. Through these descending projections, the excitability of spinal dorsal horn neurons is inhibited (Vanegas and Schaible, 2004). Decreased activity in the descending pain control systems, termed 'disinhibition', has been considered to generate a persistent pain after nerve injury (Birklein, 2002). It has been reported that activation of PAG, particular ventrolateral PAG (vlPAG), by electrical stimulation or by injection of opioids exerts analgesic action through activation of descending pain control system.

c-Fos protein, the product of the immediate early gene, is rapidly expressed in neurons in response to various stimuli, and c-Fos expression is recognized as a marker of increased neuronal activity (Lee et al., 2003). In many studies, upregulation of c-Fos expression in the vlPAG, NRM, and dorsal raphe nucleus (DR) has been suggested as the activation of descending pain control system (de Medeiros et al., 2003; Hattori et al., 2004).

Characteristic gait changes occur after unilateral sciatic nerve injury in rats. Sciatic nerve lesions cause variable loss of both extensors and flexors of the foot. This deficit causes dropping of foot to the ground and thus induces changes of the footprints. In contrast, gradual disappearance of these changes reflects nerve regeneration and functional recovery (Bervar, 2000). In this way, footprints have been used to assess the sciatic nerve function. The current and standard method for measuring functional recovery after sciatic nerve injury in rats is the sciatic function index (SFI) established by De Medinaceli et al. (1982), and subsequently modified by Bain et al. (1989). SFI formula is based on the characteristic walking patterns following sciatic nerve injury

in rats, and the recovery rate can be determined by this gait analysis.

Herbs are annual, biennial, or perennial seed-producing soft-stem plants that exhibit medicinal or aromatic properties. The use of herbal medicine as one of alternative therapies is increasing rapidly throughout in the world, which has already been used as traditional medicine in Asian countries (Tindle et al., 2005). *Harpagophytum procumbens* (Pedaliaceae), popularly known as "devil's claws", is a medicinal plant originated from Southern Africa. The plant has been traditionally used by San bushmen for a digestive tonic, headache, fever and allergy, and as an ointment to alleviate pain during childbirth. Recent clinical trials have established that *Harpagophytum procumbens* has anti-inflammatory and anti-arthritic properties for patients with degenerative joint disease (Setty and Sigal, 2005).

Atractylodes japonica (Compositae) is a perennial herb distributed in East Asia. The dried rhizomes of the plants are generally used as main ingredients in various herbal formulations for the treatment of gastrointestinal diseases (Chen et al., 2001). Aqueous extract from *Atractylodes japonica* has been reported to possess anti-inflammatory and analgesic activities through inhibition on lipopolysaccharide-induced cyclooxygenase-2 and inducible nitric oxide synthase expressions (Jang et al., 2004).

Corydalis tuber (Papaveraceae) is the root of *Corydalis yabusuo* W.T. Wang. Aqueous extract from *Corydalis tuber* has been used as an analgesic and for the treatment of inflammatory and allergic diseases (Sagare et al., 2000). The analgesic action of *Corydalis tuber* is closely associated with descending pain control system (Cheong et al., 2004).

Recently, we have developed the herbal bath named "HAC" for pain control using *Harpagophytum procumbens* extract, *Atractylodes japonica* extract, and *Corydalis tuber* extract. In the present study, the effects of herbal bath immersion on recovery rate of locomotor function and the expression of c-Fos in the various PAG regions following sciatic crushed nerve injury in rats were investigated using walking track analysis and immunohistochemistry for c-Fos.

MATERIALS AND METHODS

Experimental animals

Male Sprague-Dawley rats weighing 200 ± 10 g (6 weeks of age) were used. The experimental procedures were performed in accordance with the animal care guidelines of National Institute of Health (NIH) and the Korean Academy of Medical Sciences. The animals were housed at a controlled temperature ($20 \pm 2^\circ\text{C}$) and maintained under light-dark cycles, each consisting of 12 h of light and 12 h of darkness (lights on from 07:00 to 19:00 h), with food and water made available *ad libitum*. The rats were randomly divided into four groups ($n=8$ in each group): the sham operation group, the operation (sciatic crushed nerve injury) group, the operation and water bath group, and the operation and herbal bath group.

Surgical procedure

To induce crush injury on the sciatic nerve in rats, a surgical procedure based on previously described method was performed (De Koning et al., 1986). In brief, the right sciatic nerve was exposed through splitting incision on the gluteal muscle under pentobarbital anesthesia (50 mg/kg, i.p.; Sigma Chemical Co., St. Louis, MO, USA). The sciatic nerve was carefully exposed and crushed for 30 sec using a surgical clip between the sciatic notch and the point of trifurcation. Subsequently, the surgical wound was sutured and recovered. In the sham operation rats, the sciatic nerve was exposed but crushing pressure on the nerve was not applied.

Herbal bath immersion

After 72 h from the operation, the rats in the operation and herbal bath group were made to place in the plastic cage of 50 cm in height and 30 cm in diameter filled with herbal bath "HAC" maintained at 32°C . Herbal bath immersion was applied to the level of rat's xiphoid process for 30 min once a day for 12 consecutive days. Herbal bath "HAC" was composed of aqueous extracts of *Harpagophytum procumbens* (3.34 g/l), *Atractylodes japonica* (3.34 g/l), and *Corydalis tuber* (3.34 g/l).

The rats in the operation and water bath group were made to place in the water bath without her-

bal mixture as the same way with the rats in the operation and herbal bath group.

Walking track analysis

Functional recovery rate after sciatic nerve injury was analyzed using a walking track assessment, which can be quantified with SFI. Examination of the walking patterns was performed seven times at one day intervals through the course of the experiment as a previously described method (Bain et al., 1989). Footprints were recorded in a wooden walking alley (8.2×42 cm) with a darkened goal box at the end. The floor of the alley was covered with white paper. The anatomical landmarks on the hind feet of the rats were smeared with finger paint. The rats were allowed to walk down the track, leaving their footprints on the paper.

From the footprints, the following parameters were calculated: distance from the heel to the top of the third toe (Print Length; PL), distance between the first and the fifth toe (Toe Spread; TS), and distance from the second to the fourth toe (Intermediary Toe Spread; IT). These parameters were taken both from the intact left (non-operated) foot (NPL, NTS, and NIT) and from the injured right (experimental) foot (EPL, ETS, and EIT). SFI values were obtained using following equation (Fig. 1). Interpolating identical values of PL, TS, and IT from the right and the left hind feet are close to zero in normal rats. A value of -100 indicates complete impairment of walking ability.

c-Fos immunohistochemistry

For immunolabeling of c-Fos in the vIPAG of each brain, c-Fos immunohistochemistry was performed as a previously described method (Lee et al., 2003). Free-floating tissue sections were incubated overnight with rabbit anti-Fos antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA) at a dilution of 1 : 1,000, and the sections were then incubated for 1 h with biotinylated anti-rabbit secondary antibody (Vector Laboratories, Burlingame, CA, USA). The sections were subsequently incubated with avidin-biotin-peroxidase complex (Vector Laboratories) for 1 h at room temperature. Immunoreactivity was visualized by incubating the sections in a solution consisting of 0.05% 3,3-diaminobenzidine (DAB) and 0.01% H_2O_2 in 50 mM Tris-buffer

$$\begin{aligned} \text{SFI} = & -38.3 \left(\frac{\text{EPL} - \text{NPL}}{\text{NPL}} \right) \\ & + 109.5 \left(\frac{\text{ETS} - \text{NTS}}{\text{NTS}} \right) \\ & + 13.3 \left(\frac{\text{EIT} - \text{NIT}}{\text{NIT}} \right) \end{aligned}$$




Fig. 1. Walking track analysis. After sciatic crushed nerve injury in rats, paired parameters of the print length (PL), toe spread (TS), and intermediary toe spread (IT) were taken, and these were incorporated into Bain's formula (1989). (E) experimental side, (N) normal side, (EPL) experimental print length, (NPL) normal print length, (ETS) experimental toe spread, (NTS) normal toe spread, (EIT) experimental intermediary toe spread, (NIT) normal intermediary toe spread, (SFI) sciatic functional index.

(pH 7.6) for approximately 3 min. The sections were then washed three times with PBS and mounted onto gelatine-coated slides. The slides were air-dried overnight at room temperature, and coverslips were mounted using Permount®. As the negative control, the brain sections were likewise processed using normal goat serum in place of the primary antibody: no c-Fos-like immunoreactivity was observed.

Schematic illustration of vIPAG, chosen for the quantification of the number of Fos-positive cells is shown in Fig. 2.

Data analyses

The data are expressed as the mean±standard error mean (S.E.M). For comparisons among the groups, one-way ANOVA and Duncan's post-hoc test were performed with $p < 0.05$ as an indication of statistical significance.




Fig. 2. Schematic illustrations of the ventrolateral periaqueductal gray (vIPAG) region where the number of Fos-positive cells was counted. Aq: aqueduct, vIPAG: ventrolateral periaqueductal gray.

RESULTS

Immersion into the herbal bath enhanced SFI following sciatic crushed nerve injury

The mean SFI in each group was calculated on the 3rd, 5th, 7th, 9th, and 11th day after sciatic crushed nerve injury.

The SFI in the sham operation group was -15.17 ± 2.96 on the 3rd day, -9.73 ± 5.10 on the 5th day, -12.34 ± 3.86 on the 7th day, -18.75 ± 1.39 on the 9th day, and -16.60 ± 2.58 on the 11th day at the commencement of the experiment.

The SFI in the operation group was -98.21 ± 1.57 on the 3rd day, -97.27 ± 4.97 on the 5th day, -99.46 ± 7.64 on the 7th day, -89.61 ± 3.59 on the 9th day, and -81.06 ± 5.45 on the 11th day at the commencement of the experiment.

The SFI in the operation and water bath group was -96.76 ± 2.78 on the 3rd day, -101.93 ± 5.04 on the 5th day, -97.44 ± 6.21 on the 7th day, -89.73 ± 5.72 on the 9th day, and -88.24 ± 2.55 on the 11th day at the commencement of the experiment.

The SFI in the operation and herbal bath group




Fig. 3. Effect of herbal bath "HAC" on the sciatic functional index (SFI). The values are represented as the mean \pm S.E.M. *represents $p < 0.05$ compared to the sham operation group. (△) Sham operation group, (○) operation group, (□) operation and water bath group, and (●) operation and herb bath group.

was -97.14 ± 3.81 on the 3rd day, -95.23 ± 3.01 on the 5th day, -79.71 ± 3.71 on the 7th day, -63.59 ± 12.97 on the 9th day, and -61.36 ± 14.47 on the 11th day at the commencement of the experiment (Fig. 3).

In the present results, the SFI of the sham operation group continued near zero level during the experiment. At the beginning, the SFI in the all operation groups dropped near to -100 . In the operation group and in the operation and water bath group, the SFI value was continued at the low level until 7th day after injury and then slowly increased. In the operation and herbal bath group, SFI value was enhanced from the 7th day and rapidly increased throughout the experiment. These results indicate that immersion into the herbal bath "HAC" promotes functional locomotor recovery following sciatic crushed nerve injury.

Immersion into the herbal bath enhanced c-Fos expression in the vIPAG following sciatic crushed nerve injury

The expression of c-Fos in the vIPAG in each group was measured immediately after determination of last SFI.

The number of Fos-positive cells in the vIPAG was $71.41 \pm 3.74/\text{mm}^2$ in the sham operation group, $61.85 \pm 3.97/\text{mm}^2$ in the operation group, $63.00 \pm$




Fig. 4. Effect of herbal bath "HAC" on the c-Fos expression in ventrolateral periaqueductal gray (vIPAG). Upper: Photomicrographs of the c-Fos-positive cells. The scale bar represents 100 μm . Lower: Mean number of c-Fos-positive cells in each group. The values are represented as the mean \pm S.E.M. *represents $p < 0.05$ compared to the sham-operation group. #represents $p < 0.05$ compared to the operation group. (A) Sham operation group, (B) operation group, (C) operation and water bath group, (D) operation and herb bath group.

$4.36/\text{mm}^2$ in the operation and water bath group, and $77.80 \pm 3.77/\text{mm}^2$ in the operation and herbal bath group (Fig. 4).

In the present results, c-Fos expression in the vIPAG was reduced by sciatic crushed nerve injury and immersion into the herbal bath "HAC" significantly enhanced c-Fos expression.

DISCUSSION

Crush injury on the sciatic nerve serves as an animal model of unilateral peripheral neuropathy. Many changes affecting on the ascending facilitatory system and on the descending inhibitory system occur within the CNS as a result of neuropathy, resulting in the development of a persistent

pain. Treatment goals generally target alleviating of pain and improving of physical function (Irving et al., 2004).

The mechanisms underlying the generation of pain after peripheral nerve injury are complex involving various peripheral and central components of nervous systems and are not clarified clearly. Recent studies have proposed that the inhibition of descending pain control system caused by decreased activation of neurons is one of the mechanisms of pain production following nerve injury (Birklein, 2002; Vanegas and Schaible, 2004). Basbaum and Fields (1984) reported that electrical stimulation on the several brain stem areas elicits antinociceptive processes through activation of descending pain control system and Coimbra et al. (1992) also demonstrated that electrical or chemical stimulation into vlPAG inhibits responses to noxious stimuli.

Expression of c-Fos is commonly used to represent activation of neurons in the brain by external inputs. Upregulation of c-Fos in vlPAG, NRM, and DR induced by electroacupuncture and drugs such as morphine, antidepressant, and NMDA antagonist is associated with analgesic effect (de Medeiros et al., 2003; Hattori et al., 2004).

In the present study, c-Fos expression in the vlPAG was suppressed following sciatic crushed nerve injury, indicating decreased activity in the neurons of vlPAG. Immersion into the herbal bath "HAC" significantly enhanced c-Fos expression in the vlPAG. The present results show that herbal bath "HAC" facilitates neuronal activity in the vlPAG following sciatic nerve injury.

The analgesic effect of several kinds of herbal extract on the neuropathic pain has been well reported. Tatsumi et al. (2004) demonstrated that extracts of *Moutan cortex* and *Coicis semen* have an analgesic effect on the neuropathic pain in mice. Analgesic effect of certain herbs has been suggested to be involved in the descending pain control system. Isono et al. (1994) and Omiya et al. (1999) showed that antinociceptive action of *Aconiti tuber* is implicated in the descending pain control system. Shin et al. (2003) demonstrated that *Chelidonii herba* increases neuronal excitability in PAG, which results in activation of descending pain control system and may contribute as a potential me-

chanism of the analgesic actions of *Chelidonii herba*. Cheong et al. (2004) also reported that application of *Corydalis tuber* onto PAG neurons modulates glycine-activated ion current in the PAG neurons, which exerts analgesic action.

The SFI derived from walking track analysis in rats provides a reliable and easily quantifiable method for the assessing of motor function after sciatic nerve injury (Varejão et al., 2003). This gait analysis is based on the fact that rats normally walk on their digits and metatarsal footpads. Print length is therefore short in normal animals. Sciatic nerve injury causes functional loss of both extensor muscles and flexor muscles of the foot, causing drop of foot.

In sciatic nerve crushed injury model, Vogelaar et al. (2004) reported that although sensory and motor reinnervation of the paw are fully established after nerve injury, persistent pain still exists and the animals can not support their weight on the injured paw. In the acute stage of sciatic crushed nerve injury, we observed that the flexion contracture of the toes and a curvation of the feet make impossible to calculate SFI in some rats. The rats subjected to crush injury sometimes walk by their dorsum of the affected foot or load their weight on the medial part of affected foot. These observations may be due to compensatory immobilization to painful dysesthesia as well as neurological loss.

In the present study, right sciatic crushed nerve injury in rats resulted in the characteristic pattern of the footprints, representing reduction in the SFI value. The SFI value of the rats in the operation and herb bath group was significantly increased from 7th day of the experiment, whereas the SFI value of the rats in the operation group and in the operation and water bath group remained low level until 11th day of the experiment. The present results indicate that herbal bath "HAC" accelerates functional recovery from the locomotor deficit after sciatic crushed nerve injury. The present study implies that decreased activation of descending pain control system induced by sciatic nerve injury may consequently contribute muscle atrophy and motor dysfunction. Herbal bath "HAC" may facilitate motor performance through stimulation of the descending pain control system by activating neurons in the vlPAG.

Here in this study, we have shown that herbal bath "HAC" activates neurons in the vIPAG, and thereby facilitates motor function. Based on the present results, herbal bath "HAC" can be used as a new therapeutic intervention for pain control and functional recovery from peripheral nerve injury.

ACKNOWLEDGMENTS

This research was supported by Basic Science Research Program Through the National Research Foundation (NRF) funded by the Ministry of Education, Science and Technology (R11-2005-014).

REFERENCES

- Bain JR, Mackinnon SE and Hunter DA (1989) Functional evaluation of complete sciatic, peroneal, and posterior tibial nerve lesions in the rat. *Plast Reconstr Surg* 83:129-138.
- Basbaum AI and Fields HL (1984) Endogenous pain control systems: brainstem spinal pathways and endorphin circuitry. *Annu Rev Neurosci* 7:309-338.
- Bennett GJ and YK Xie (1988) A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. *Pain* 33:87-107.
- Bervar M (2000) Video analysis of standing-an alternative footprint analysis to assess functional loss following injury to the rat sciatic nerve. *J Neurosci Methods* 102:109-116.
- Birklein F (2002) Mechanisms of neuropathic pain and their importance in Fabry disease. *Acta Paediatr Suppl* 91:34-37.
- Chen KT, Su YC, Lin JG, Hsin LH, Su YP, Su CH, Li SY, Cheng JH and Mao SJ (2001) Identification of Atractylodes plants in Chinese herbs and formulations by random amplified polymorphic DNA. *Acta Pharmacol Sin* 22:493-497.
- Cheong BS, Choi DY, Cho NH, Lee JD, Chang HK, Shin MC, Shin MS and Kim CJ (2004) Modulation of Corydalis tuber on glycine-induced ion current in acutely dissociated rat periaqueductal gray neurons. *Biol Pharm Bull* 27:1207-1211.
- Coimbra NC, Tomaz C and Brandão ML (1992) Evidence for the involvement of serotonin in the antinociception induced by electrical or chemical stimulation of the mesencephalic tectum. *Behav Brain Res* 50:77-83.
- De Koning P, Brakkee JH and Gispen WH (1986) Methods for producing a reproducible crush in the sciatic and tibial nerve of the rat and rapid and precise testing of return of sensory function. Beneficial effects of melanocortins. *J Neurol Sci* 74:237-246.
- De Medeiros MA, Canteras NS, Suchecki D and Mello LE (2003) Analgesia and c-Fos expression in the periaqueductal gray induced by electroacupuncture at the Zusanli point in rats. *Brain Res* 973:196-204.
- De Medinaceli L, Freed WJ and Wyatt RJ (1982) An index of the functional condition of rat sciatic nerve based on measurements made from walking tracks. *Exp Neurol* 77:634-643.
- Hattori Y, Watanabe M, Iwabe T, Tanaka E, Nishi M, Aoyama J, Satoda T, Uchida T and Tanne K (2004) Administration of MK-801 decreases c-Fos expression in the trigeminal sensory nuclear complex but increases it in the midbrain during experimental movement of rat molars. *Brain Res* 1021:183-191.
- Irving G, Goli V and Duntzman E (2004) Novel pharmacologic options in the treatment of neuropathic pain. *CNS Spectr* 9:1-11.
- Isono T, Oyama T, Asami A, Suzuki Y, Hayakawa Y, Ikeda Y, Noguchi M and Omiya Y (1994) The analgesic mechanism of processed Aconiti tuber: the involvement of descending inhibitory system. *Am J Chin Med* 22:83-94.
- Jang MH, Lim S, Han SM, Park HJ, Shin I, Kim JW, Kim NJ, Lee JS, Kim KA and Kim CJ (2003) Harpagophytum procumbens suppresses lipopolysaccharide-stimulated expressions of cyclooxygenase-2 and inducible nitric oxide synthase in fibroblast cell line L929. *J Pharmacol Sci* 93:367-371.
- Jang MH, Shin MC, Kim YJ, Kim CJ, Kim Y and Kim EH (2004) Atractylodes japonica suppresses lipopolysaccharide-stimulated expressions of inducible nitric oxide synthase and cyclooxygenase-2 in RAW 264.7 macrophages. *Biol Pharm Bull* 27:324-327.
- Lee MH, Kim H, Lim BV, Chang HK, Lee TH, Jang MH, Shin MC, Lee J, Shin MS and Kim CJ (2003) Naloxone potentiates treadmill running-induced increase in c-Fos expression in rat hippocampus. *Life Sci* 73:3139-3147.
- Omiya Y, Goto K, Suzuki Y, Ishige A and Komatsu Y (1999) Analgesia-producing mechanism of processed Aconiti tuber: role of dynorphin, an endogenous kappa-opioid ligand, in the rodent spinal cord. *Jpn J Pharmacol* 79:295-301.
- Sagare AP, Lee YL, Lin TC, Chen CC and Tsay HS (2000) Cytokinin-induced somatic embryogenesis and plant regeneration in Corydalis yanhusuo (Fumariaceae) - a medicinal plant. *Plant Sci* 160:139-147.
- Setty AR and Sigal LH (2005) Herbal medications commonly used in the practice of rheumatology: mechanisms of action, efficacy, and side effects. *Semin Arthritis Rheum* 34:773-784.
- Shin MC, Jang MH, Chang HK, Lim S, Han SM, Park HJ, Shim I, Lee JS, Kim KA and Kim CJ (2003) Modulation of Chelidonium herba on glycine-activated and glutamate-activated ion currents in rat periaqueductal gray neurons. *Clin Chim Acta* 337:93-101.
- Tatsumi S, Mabuchi T, Abe T, Xu L, Minami T and Ito S (2004) Analgesic effect of extracts of Chinese medicinal herbs Moutan cortex and Coicis semen on neuropathic pain in mice. *Neurosci Lett* 370:130-134.
- Tindle HA, Davis RB, Phillips RS and Eisenberg DM (2005) Trends in use of complementary and alternative medicine by US adults: 1997-2002. *Altern Ther Health Med* 11:42-49.
- Vanegas H and Schaible HG (2004) Descending control of persistent pain: inhibitory or facilitatory? *Brain Res Brain Res Rev* 46:295-309.
- Varejão AS, Cabrita AM, Geuna S, Melo-Pinto P, Filipe VM, Gramsbergen A and Meek MF (2003) Toe out angle: a

functional index for the evaluation of sciatic nerve recovery in the rat model. *Exp Neurol* 183:695-699.

Vogelaar CF, Vrinten DH, Hoekman MF, Brakkee JH, Burbach JP and Hamers FP (2004) Sciatic nerve regeneration in

mice and rats: recovery of sensory innervation is followed by a slowly retreating neuropathic pain-like syndrome. *Brain Res* 1027:67-72.